

Biovit- M

Multivitamin with minerals Tablet

Presentation:

Each sugar free film coated tablet contains :

Vitamin A Acetate	USP	1.5 mg
Vitamin D ₃	USP	10.0 mcg
Thiamine Mononitrate	USP	1.5 mg
Riboflavin	USP	1.7 mg
Pyridoxine Hydrochloride	USP	2.0mg
Nicotinamide	USP	20.0 mg
Ascorbic Acid	USP	60.0 mg
Folic Acid	USP	0.4 mg
Calcium Pantothenate	USP	10.92 mg
Cyanocobalamin	USP	6.0 mcg
Vitamin E Acetate	BP	15.0 mg
Dried Ferrous Sulphate	BP	50.0mg
Potassium Iodide	USP	196 mcg
Potassium Sulphate	BP	11.141 mg
Manganese Sulphate Monohydrate	BP	1.0 mg
Cupric Sulphate Pentahydrate	BP	2.0 mg
Zinc Sulphate Monohydrate	BP	37.03mg

Indications:

Deficiency states in acute and chronic disease. Conditions regarding specific support : pregnancy, lactation, menopause, during treatment with antibiotics. The comprehensive formulae of Biovit- M assures liberal amounts of important vitamins, minerals and trace elements needed by the body during periods of increased energy requirements such as in disease and convalescence.

Dosage and Administration:

One tablet daily or as advised by the physician.

Contraindications:

Supplemental vitamins and minerals should not be prescribed for patients with haemochromatosis or Wilson's disease. Hypersensitivity to any of the active ingredients is a contraindication. Excessive doses of vitamin A and D can lead to hypervitaminosis. When multivitamin preparations are prescribed allowance must be made for vitamins from other sources. This is particularly important during the first trimester of pregnancy when large doses of Vitamin A may be teratogenic. Iron and Zinc chelate with tetracyclines and absorption of all three agents may be impaired.

Precautions:

Biovit-M may be used in patients being treated for Parkinson's disease with a Levodopa preparation which includes a dopa-decarboxylase inhibitor (e.g., Carbidopa, Benserazide) but precautions should be taken. Pyridoxine acts as an antagonist of Levodopa.

Side Effects:

Multivitamin preparation with ordinary doses of component are usually nontoxic.

Use in pregnancy & lactation:

Recommended.

Drug interactions:

No such drug interactions have been reported.

Storage Condition:

Store at 25⁰ C in a dry place. Keep away from light.

Warning:

To be sold and dispensed only on the prescription of a registered physician. Keep out of reach of children. Excessive dosage of Vitamin A would be dangerous to the health. Before long term use please consult your doctor.

Commercial Pack:

Biovit-M Tablet : Amber Glass Bottle containing 30 sugar free film coated tablets.

Manufactured by



For further query on the use of this medicine, consult to a registered Doctor or Pharmacist.