

Neurep

Vitamin B1, B6 & B12

Composition:

Each film coated tablet contains:

Neurep Tablet: Thiamine Mononitrate (Vitamin B1) BP : 100 mg
Pyridoxine Hydrochloride (Vitamin B6) BP : 200 mg
Cyanocobalamin (Vitamin B12) BP : 200 mcg

Description:

Neurep is a combination of Vitamin B1 (Thiamine Mononitrate), B6 (Pyridoxine Hydrochloride) & B12 (Cyanocobalamin). These vitamins play essential role as co-enzyme for the metabolism of nervous system. Thus this combination normalizes the nerve cell metabolism. Neurep supports the regeneration of nerve fibers and myelin sheath by activation of the metabolism and the natural repair mechanism. The combination of these 3 vitamins shows better efficacy than that of the individual components. In addition, Vitamin B1, B6 & B12 promote body's own repair mechanism and work in pain management.

Indications:

Chronic pains like sciatica, lumbago, trigeminal neuralgia, facial paralysis, optic neuritis as well as neuropathy & cardiac complications. Besides these, deficiency exists of relevant vitamins.

Dosage and administration:

Neurep Tablet: 1-2 tablets three times daily.

Side-effects:

Very rarely hypersensitive reaction and sensitivity disturbance may occur.

Use in pregnancy:

Neurep is safe for pregnant woman.

Use in lactating mother:

It is not known whether Thiamine Mononitrate (Vitamin B1) is excreted in breast milk or not, but may be taken cautiously during lactation. Vitamin B6 & B12 have no adverse effect during lactation.

Children: Neurep may be given to children.

Precautions:

This combination (Vitamin B1+B6+B12) should be avoided in case of hypersensitivity to Thiamine Mononitrate, cobalt or any other component of these ingredients. Cyanocobalamin should not be given before a diagnosis has been fully established because of the masking symptoms of subacute degeneration of the spinal cord.

Contraindication:

It should not be used in patients on levodopa therapy.

Drug Interactions:

Pyridoxine Hydrochloride (Vitamin B6) reduces therapeutic effect of levodopa but it does not occur if a dopa decarboxylase inhibitor is also given. Neomycin, chloramphenicol, aminosallylic acid & histamine H2 receptor may reduce Vitamin B12 absorption.

Storage:

For Tablet: Store in a cool and dry place, protected from light. Keep out of the reach of children.

Commercial Pack:

Neurep Tablet: Each box contains 6x10's Neurep Tablets in blister pack.

Manufactured by



For further query on the use of this medicine, consult to a registered Doctor or Pharmacist.