

I-fort

Polyethylene Glycol USP 0.4% and
Propylene Glycol USP 0.3%
Sterile Eye Drops

Composition

I-fort Sterile Eye Drops: Each ml contains Polyethylene Glycol 400 USP 4 mg & Propylene Glycol USP 3 mg.

Clinical Pharmacology

The combination of Polyethylene Glycol 400 & Propylene Glycol is clinically proven to reduce both signs and symptoms of dry eye. The mechanism of action is thought to be due to its unique gelling and lubricating system formulated to adjust to each users individual tear pH. When the ingredients of I-fort combine with natural tears, a soft gel forms a network of protection over the eye surface. Since it promotes a healthy environment in eye surface, damaged surface cells of eye can repair more easily.

Indications and Uses

I-fort Sterile Eye Drops is indicated for the temporary relief of burning and irritation due to dryness of the eye.

Dosage and Administration

Instill 1 drop 4 times daily in the affected eye(s) or as needed.

Contraindications

It is contraindicated in patients with known hypersensitivity to any ingredient in this formulation.

Warning & Precautions

- Never touch tip of container with any surface to avoid contamination.
- Replace cap after each use.

Side Effects

Generally well tolerated. I-fort should not be used if allergic condition occurs to any ingredients of the product.

Storage

Store in a cool and dry place, protect from light. Do not use longer than one month after first opening. Keep out of the reach of children.

Commercial Pack

I-fort Sterile Eye Drops: Each plastic dropper bottle contains 10 ml sterile eye drops.

Manufactured by

General Pharmaceuticals Ltd.
Kaliakoir, Gazipur for



For further query on the use of this medicine, consult to a registered Doctor or Pharmacist.